

Tagging the $pn \rightarrow d\phi$ reaction by backward protons in $pd \rightarrow d\phi p_{sp}$ processes

L.P. KAPTARI^{a,b,c}, B. KÄMPFER^a, S.S. SEMIKH^{a,b}

^a *Forschungszentrum Rossendorf, PF 510119, 01314 Dresden, Germany*

^b *Bogolyubov Laboratory of Theoretical Physics, JINR, Dubna 141980, Russia*

^c *Department of Physics, University of Perugia, and INFN Sezione di Perugia, via A. Pascoli, I-06100, Italy*

Abstract

The reaction $pd \rightarrow d\phi p_{sp}$ is studied within the Bethe-Salpeter formalism. Under special kinematical conditions (slow backward spectator proton p_{sp} and fast forward deuteron) relevant for forthcoming experiments at COSY, the cross section and a set of polarization observables factorize in the contribution of the pure subprocess $pn \rightarrow d\phi$ and a contribution stemming from deuteron quantities and kinematical factors. This provides a theoretical basis for studying threshold-near processes at quasi-free neutrons.

PACS: 13.75-n, 14.20.-c, 21.45+v

Keywords: hadron-induced intermediate-energy reaction, meson production, few-body systems

I. INTRODUCTION

Data of elementary nucleon reactions involving a neutron in the entrance channel are scarce. One can use, for instance, a tagged neutron (n) beam [1] impinging on a proton (p) target or a proton beam impinging on a nuclear target. In the latter case one needs models to deconvolute the pn reaction. Here we are interested in threshold-near meson production. Over the last decade the data basis for the pp reactions has been impressively improved [2]. Such a detailed knowledge of elementary hadron reactions is necessary for analyzing heavy-ion collisions. Threshold-near meson production is also interesting for testing predictions of QCD in the non-perturbative region, provided e.g., by chiral perturbation theory. While it seems to be straightforward to translate cross sections for reactions of the type $pp \rightarrow ppM$ (M is a meson) into the cross section $nn \rightarrow nnM$, the channel $pn \rightarrow pnM$ is less simple accessible. Model calculations point to a non-trivial energy dependence of the ratio $\sigma(pp \rightarrow ppM)/\sigma(np \rightarrow npM)$, e.g. for ϕ meson production [3]. Experiments showed, furthermore, a surprisingly large ratio $\sigma(pd \rightarrow pd\eta)/\sigma(pp \rightarrow pp\eta)$ [1,4]. Therefore, it is desirable to test directly predictions for the pn reactions.

The spectator technique [5,6] offers one possibility to isolate to a large extent the quasi-free reaction at a neutron when a proton beam impinges on a deuteron target. Since the final state interactions in the pn system differs for singlet and triplet states and can be large

at threshold, it is of great advantage to consider the particular reaction $pn \rightarrow dM$, as the deuteron d is in one well defined state. Guided by this, in [7,8] the reactions $pn \rightarrow dM$ with $M = \eta, \eta', \omega, \phi, a_0^+$ have been considered.

The aim of the present note is to analyze the complete reaction $pd \rightarrow dMp_{sp}$ and to test to what extent the spectator technique [5,6] really selects a quasi-free reaction at the neutron in the deuteron target. In other words, we are going to derive a factorization theorem showing kinematical conditions for which the total cross section $\sigma(pd \rightarrow dMp_{sp})$ factorizes in a part depending on the target deuteron characteristics and a part $\sigma(pn \rightarrow dM)$. In doing so we select as a sufficiently transparent example the ϕ production. The reasons are obvious: (i) the elementary ϕ production amplitude is simple [3], (ii) the ϕ meson is interesting with respect to OZI rule [9] and hidden strangeness in the nucleon [3,8,10,11], and (iii) the ϕ meson is important with respect to both the inclusive K^- production [12] and the e^+e^- decay channel to be studied at HADES in near future [13]. An extension to ρ and ω production goes along the same lines, but due to the role of s, u channel baryon resonances [10,14,15] the elementary amplitude of the subprocess $pn \rightarrow pnM$ is by far more complicate and needs separate consideration, as other mesons too.

Our paper is organized as follows. In section II we present the kinematics, cross section and amplitudes of the complete reaction $pd \rightarrow d\phi p_{sp}$. Numerical results are exhibited in section III, where polarization observables are discussed and compared with the ones of the quasi-free sub-reaction $pn \rightarrow d\phi$. We also derive selection rules. The factorization of the cross section is considered in section IV. The summary can be found in section V.

II. KINEMATICS, CROSS SECTION, AMPLITUDES

Let us consider the process of ϕ meson production in the exclusive reaction envisaged in [6]

$$p_p + p_d = p'_d + p_\phi + p'_p, \quad (1)$$

at kinetic energies of the incoming proton corresponding to those achievable at the Cooler Synchrotron COSY $T_p \leq 2.7$ GeV. The deuteron is here supposed to be detected in the forward direction, i.e., in direction of the incoming proton with relativistic energy, $T'_d \sim 1$ GeV, and, contrarily, the final (spectator) proton is slowly moving in the backward direction ($\theta_{sp} > 90^\circ$, $p'_p < 100$ MeV). The main contribution to the process (1) comes then from the spectator mechanism, where the incoming proton interacts with the internal neutron from the deuteron and produces a meson and a deuteron in the final state. The second proton in the deuteron acts merely as a spectator. Schematically the spectator mechanism is depicted in Fig. 1.

In the laboratory system, where the target deuteron is at rest, the relevant momenta are defined by $p_p = (E_p, \mathbf{p}_p)$ and $p_d = (M_d, \mathbf{0})$ as initial momenta of the proton and deuteron, $p'_p = (E'_p, \mathbf{p}'_p)$, $p'_d = (E'_d, \mathbf{p}'_d)$, and $p_\phi = (E_\phi, \mathbf{p}_\phi)$ as momenta of final proton, deuteron and ϕ meson, respectively. The fivefold differential cross section reads

$$2E'_d \frac{d^5\sigma}{d^3p'_d d\Omega'_p} = \frac{1}{8(2\pi)^5\sqrt{\lambda}} \left[\frac{|\mathbf{p}'_p|^2}{|B|\mathbf{p}'_p| - CE'_p|} \right] \frac{1}{6} \sum_{spins} |\mathcal{F}|^2, \quad (2)$$

where the Källén function $\lambda = 4 M_d^2 |\mathbf{p}_p|^2$ and the kinematical coefficients $B = M_d + E_p - E'_d$ and $C = |\mathbf{p}_p| \cos \theta'_p - |\mathbf{p}'_d| \cos \theta'_{dp}$ reflect the energy conservation. The invariant amplitude $\mathcal{F}$ depends, besides the spin projections $s_p, \mathcal{M}_d; s'_p, \mathcal{M}'_d, \mathcal{M}_\phi$ on the quantization z-axis (oriented along $\mathbf{p}_p$), also upon the initial energy and five independent kinematical variables, say $\mathbf{p}'_d$ and Ω'_p .

To compute the amplitude $\mathcal{F}$, let us define the scattering operator $\hat{\mathcal{O}}_{\mathcal{M}_\phi, \mathcal{M}'_d}$ (a $4 \otimes 4$ matrix in the spinor space) as the truncated amplitude for the subprocess $pn \rightarrow d\phi$ (hatched blob of the diagram in Fig. 1), i.e., as an operator which, when sandwiched between two free spinors u_p and u_n , results in the amplitude of the real process $pn \rightarrow d\phi$ with two on-mass shell nucleons. In the general case this operator is assumed to describe the process of meson production from two off-mass shell nucleons. With this definition, the amplitude $\mathcal{F}$ may be written as

$$\mathcal{F} = \bar{u}_\alpha(p'_p, s'_p) (\hat{p}'_p - m)_{\alpha\beta} \Phi_{\mathcal{M}_d}^{c\beta}(p_n, p'_p) \hat{\mathcal{O}}_{\mathcal{M}_\phi, \mathcal{M}'_d}^{\delta c} u_\delta(p_p, s_p), \quad (3)$$

where m is the nucleon mass and $\Phi_{\mathcal{M}_d}^{c\beta}(p_n, p'_p)$ denotes the Bethe-Salpeter (BS) amplitude of the deuteron in the momentum space. In Eq. (3), the spinor indices are exhibited explicitly, with Latin and Greek characters referring to neutrons and protons, respectively (summation over indices occurring pairwise is supposed). An expression for the amplitude (3) can be found by specifying a model for the operator $\hat{\mathcal{O}}$. However, in order to keep the correspondence between the reaction (1) and the process of ϕ meson production in NN collisions as close as possible, we highlight explicitly its dependence upon the BS amplitude of the final deuteron by introducing another operator $\hat{\mathcal{O}}$ as

$$\hat{\mathcal{O}}_{\mathcal{M}_\phi, \mathcal{M}'_d}^{\delta c} = -i \int \frac{d^4 p}{(2\pi)^4} \bar{\Phi}_{\mathcal{M}'_d}^{\alpha b}(1', 2') \hat{\mathcal{O}}_{\alpha\delta}^{bc}(12; 1'2', \mathcal{M}_\phi). \quad (4)$$

Then from Eq. (4) and from Fig. 1 it is easy to recognize the operator $\hat{\mathcal{O}}(12; 1'2', \mathcal{M}_\phi)$ as the one describing the ϕ meson production in the sub-reaction $pn \rightarrow d\phi$ with an off-mass shell initial neutron. Such an operator has been introduced in Ref. [16] where its properties have been discussed in detail. We briefly recall here the main steps in computing $\hat{\mathcal{O}}(12; 1'2', \mathcal{M}_\phi)$ and the amplitude $\mathcal{F}$ in Eq. (3). Usually, within the BS formalism in order to exhibit formulae in a familiar matrix form, one introduces new, “charge-conjugated” quantities such as $\Psi \equiv -\Phi\gamma_c$, $\hat{\mathcal{Q}} \equiv \gamma_c \hat{\mathcal{O}}$, $\gamma_c \gamma_\mu^T = -\gamma_\mu \gamma_c$, $v(p) = [\bar{u}(p)\gamma_c]^T$, $\bar{v}(p) = [\gamma_c u(p)]^T$. Then the amplitude $\mathcal{F}$ can be written as

$$\mathcal{F} = \bar{v}(p_p, s_p) \hat{\mathcal{Q}}_{\mathcal{M}_\phi, \mathcal{M}'_d} \Psi_{\mathcal{M}_d}(p_n, p'_p) (\hat{p}'_p + m) v(p'_p, s'_p), \quad (5)$$

where in Eq. (5) and in subsequent expressions one has usual matrix multiplications and traces. For further simplification we insert between $\hat{\mathcal{Q}}$ and Ψ in (5) the complete set of the Dirac free spinors $\hat{I} = (1/2m) \sum_{r=1}^4 \varepsilon_r u_r(p) \bar{u}_r(p)$, where $\varepsilon_r = +1$ for $r = 1, 2$ and $\varepsilon_r = -1$ for $r = 3, 4$, to obtain

$$\mathcal{F} = \frac{1}{2m} \sum_{s_n} T_{s_p, s_n}^{\mathcal{M}_\phi, \mathcal{M}'_d}(\nu_{pn}, t) \bar{u}(p_n, s_n) \Psi_{\mathcal{M}_d}(p_n, p'_p) (\hat{p}'_p + m) v(p'_p, s'_p). \quad (6)$$

The amplitude $T_{s_p, s_n}^{\mathcal{M}_\phi, \mathcal{M}'_d}(\nu_{pn}, t) = \bar{v}(p_p, s_p) \hat{\mathcal{Q}}_{\mathcal{M}_\phi, \mathcal{M}'_d} u(p_n, s_n)$ coincides with the matrix element of the real process $pn \rightarrow d\phi$, defined by the Mandelstam variables $\nu_{pn} = (p_p + p_n)^2$, $t = (p_p - p'_d)^2$. Here is worth emphasizing that the kinematics of the process (1) differs from the one of the real process $pn \rightarrow d\phi$. In spite of commonly defined variable t , the numerical values of momenta of the final meson and deuteron do not coincide in the two processes. Rather, the amplitude $T_{s_p, s_n}^{\mathcal{M}_\phi, \mathcal{M}'_d}(\nu_{pn}, t)$ in the process (1) corresponds to an amplitude of meson production from an off-mass shell neutron. If the off-mass shell effects were negligibly small, then $T_{s_p, s_n}^{\mathcal{M}_\phi, \mathcal{M}'_d}(\nu_{pn}, t)$ in both processes would depend solely upon the momentum transfer t and the effective energy $\nu_{NN} = (E_p + E'_p)^2 - (\mathbf{p} - \mathbf{p}')^2$. As shown in Ref. [16], $T_{s_p, s_n}^{\mathcal{M}_\phi, \mathcal{M}'_d}$ can be written in the form

$$T_{s_p, s_n}^{\mathcal{M}_\phi, \mathcal{M}'_d}(\nu_{pn}, t) = \frac{i}{(2m)^2} \sum_{r, r'=1}^4 \int \frac{d^4 p}{(2\pi)^4} A_{s_p, s_n; r, r'}^{\mathcal{M}_\phi}(12; 1'2') \bar{v}_{r'}(2') \bar{\Psi}_{\mathcal{M}'_d}(1', 2') u_r(1'), \quad (7)$$

where the partial spin amplitudes $A_{s_p, s_n; r, r'}^{\mathcal{M}_\phi}(12; 1'2')$ can be calculated within an effective one-boson exchange model by an explicit evaluation of the elementary diagrams depicted in Fig. 2 in [16]. We use the parameter set “B” in Ref. [11] to calculate the amplitudes $A_{s_p, s_n; r, r'}^{\mathcal{M}_\phi}(12; 1'2')$. Now the second part of the matrix element in Eq. (6), which includes the BS amplitude $\Psi_{\mathcal{M}_d}$ for the deuteron at rest, can be straightforwardly evaluated by using the explicit expressions for the ${}^3S_1^{++}$ and ${}^3D_1^{++}$ components of the deuteron BS amplitude [17]. The result is

$$\mathcal{F} = \sqrt{8\pi M_d} \sum_{s_n} T_{s_p, s_n}^{\mathcal{M}_\phi, \mathcal{M}'_d}(\nu_{pn}, t) \left[u_S(|\mathbf{p}'|) C_{\frac{1}{2}s_p \frac{1}{2}s_n}^{1\mathcal{M}_d} - u_D(|\mathbf{p}'|) \sqrt{4\pi} \sum_{m, \mu} C_{\frac{1}{2}s_p' \frac{1}{2}s_n}^{1\mu} C_{2m1\mu}^{1\mathcal{M}_d} Y_{2m}(\mathbf{n}) \right], \quad (8)$$

where M_d is the deuteron mass, $u_{S,D}$ denote S and D waves the BS wave function [17,18], $\mathbf{n} = \mathbf{p}/|\mathbf{p}|$ and $Y_{2m}(\mathbf{n})$ the spherical harmonics of second order. In deriving Eq. (8) we used the identity $\sqrt{2} \sum_{\nu_1 \nu_2} C_{\frac{1}{2}\nu_1 \frac{1}{2}\nu_2}^{1\mathcal{M}_d} \chi_{\frac{1}{2}\nu_1}^T \chi_{\frac{1}{2}\nu_2}^T = i(\boldsymbol{\sigma} \boldsymbol{\xi}_{\mathcal{M}_d}) \sigma_y$, where $C_{\frac{1}{2}\nu_1 \frac{1}{2}\nu_2}^{1\mathcal{M}_d}$ are the Clebsch-Gordan coefficients, $\chi_{\frac{1}{2}\nu}$ and $\boldsymbol{\xi}$ stand for the spin- $\frac{1}{2}$ functions and the deuteron polarization vector, respectively. Equation (8) shows that an exact factorization of the cross section into two parts (one depending upon the target deuteron wave function, another one upon the kinematics and dynamics of the subprocess $pn \rightarrow d\phi$) generally does not occur. This is because the summation over the spins of the internal neutron s_n which leads to interference terms in the square of the matrix element. However, below we show that under special kinematical conditions an approximate factorization is possible.

III. NUMERICAL RESULTS

From Eqs. (6 - 8), the differential cross section (2) and any other polarization observables of the process $pd \rightarrow d\phi p_{sp}$ are calculable. The BS wave functions $u_{S,D}$ basically coincide with the non-relativistic S and D waves of the deuteron for small values of the momentum of the spectator proton (see [17]). Therefore, the relativistic and non-relativistic expressions for the amplitude $\mathcal{F}$ have the same form, and one may perform also calculations by using

deuteron wave functions obtained within a non-relativistic approach. In our calculations we use the BS solution [18], obtained with a realistic one-boson exchange kernel. Additionally, Paris and Bonn wave functions are considered to investigate the sensitivity of observables on the deuteron model.

All our calculations have been performed within kinematical conditions corresponding to the COSY Proposal #75 [6], i.e., the detected deuteron is supposed to be in the forward direction and the spectator proton in the backward hemisphere with $\theta'_p \sim 120^\circ - 140^\circ$. The momenta of the deuteron are chosen in the range $p'_d \sim 1.9 - 3$ GeV/c, which means that the spectator proton is detected with low velocity, i.e., $p'_p \sim 0 - 0.04$ GeV/c. These kinematic conditions imply that both the reaction (1) and the subprocess $pn \rightarrow d\phi$ occur essentially near the threshold.

In Fig. 2, the differential cross section is presented as a function of the deuteron momentum for the specific choice of variables $\theta'_d = 0^\circ$, and $\theta'_p = 130^\circ$. (We changed the kinematics with θ'_d up to 5° and find a very smooth behavior of all observable.) The kinetic energy of the initial proton is $T_p = 2.69$ GeV. The maximum of the cross section at $p'_d \sim 2.35$ GeV/c corresponds to that kinematical condition where the momentum of the spectator proton is maximal, $p'_p \sim 0.035$ GeV/c. Together with the large spectator angles this implies that at these kinematical conditions the excess energy in the $pn \rightarrow d\phi$ subprocess is minimal ($\Delta s^{1/2} \sim 4$ MeV) and very near the threshold. Obviously, since at such kinematics the production amplitude is almost constant, the maximum in the differential cross section has a pure kinematical origin being entirely governed by the kinematical factor within the square brackets in Eq. (2). Fig. 2 highlights that different models for the deuteron wave function result in quantitatively different cross sections. This is because the integration over the internal momentum in the final deuteron covers the region of the minimum of the S wave where different models provide essentially different wave functions (see discussion in [16]).

Let us now discuss polarization observables. Since the spin structure of the amplitude $\mathcal{F}$ in Eq. (8) is basically determined by the spin structure of the amplitude $T_{s_p, s_n}^{\mathcal{M}_\phi, \mathcal{M}'_d}(\nu_{pn}, t)$ of the subprocess $pn \rightarrow d\phi$, we proceed with an analysis of this amplitude. Bearing in mind that $T_{s_p, s_n}^{\mathcal{M}_\phi, \mathcal{M}'_d}(\nu_{pn}, t)$ is a Lorentz invariant quantity one can, for the sake of simplicity, investigate its structure in the center of mass of the detected deuteron (near the threshold this system coincides with the center of mass system of two nucleons), provided the Lorentz boost is performed along the z -axis. In such a case the spin structure of $T_{s_p, s_n}^{\mathcal{M}_\phi, \mathcal{M}'_d}(\nu_{pn}, t)$ is the same as in the laboratory system (no additional Wigner rotations are needed). Near the threshold, the final $\phi - d$ system is in a S orbital state corresponding to $I_f = 0$, $J_D^\pi = 1^+$ ($L_D = 0, 2$), $J_f^\pi = J_\phi^\pi + J_D^\pi = 0^-, 1^-, 2^-$, where I , L and J^π are the total isospin, total radial angular momentum, angular momentum and parity, respectively. Thus from symmetry constraints the allowed configurations in the initial state are $I_i = 0$, $L_i = 1, 3, 5 \dots$ and total spin $S_i = 0$. The conservation law $J_i = J_f$ implies $L_i = 1$, so that $J_i^\pi = 1^-$. Hence, near the threshold the main contribution into the amplitude $T_{s_p, s_n}^{\mathcal{M}_\phi, \mathcal{M}'_d}(\nu_{pn}, t)$ comes from initial spin configurations with $s_p = -s_n$ and, since the two nucleons in the final state form a deuteron (with the total spin $S = 1$), the most probable transitions near the threshold are expected to be those with nucleon spin-flips in the corresponding vertices and with the deuteron projections $\mathcal{M}'_d = \pm 1$. Consequently, the projections of the ϕ meson are to be $\mathcal{M}_\phi = \mp 1$.

Coming back to the complete reaction one can expect that the prevailing contribution

into the full amplitude $\mathcal{F}$ comes from configurations with $\mathcal{M}_d - s'_p = -s_p$, if (i) the effects of D waves in the target deuteron may be disregarded near the threshold, (ii) the spectator mechanism describes the reaction appropriately, and (iii) the factorization (see below) holds. Hence, near the threshold the polarization observables of the reaction (1) are predicted to behave in the same manner as the ones of the subprocess $pn \rightarrow d\phi$. Therefore, the study of various polarization observables allows to check the factorization.

Let us define the asymmetry of the reaction as

$$\mathcal{A} = \frac{d\sigma(s_p + s_n = 1) + d\sigma(s_p + s_n = -1) - d\sigma(s_p + s_n = 0)}{d\sigma(s_p + s_n = 1) + d\sigma(s_p + s_n = -1) + d\sigma(s_p + s_n = 0)}, \quad (9)$$

where the spin of the intrinsic neutron is defined as $s_n = \mathcal{M}_d - s'_p$. Then, from the above selection rules near the threshold, the asymmetry $\mathcal{A}$ is predicted to be negative and close to -1. In Fig. 3, results of a complete calculation of the asymmetry $\mathcal{A}$ are presented. The kinematical conditions are as in Fig. 2. It is seen that in the whole kinematical range of p'_d the asymmetry is basically the same as in the subprocess $pn \rightarrow d\phi$. Moreover, as expected, at p'_d near the maximum the asymmetry approaches -1.

Similar results are obtained for other polarization observables. For instance, the tensor analyzing power defined as

$$T_{20} = \frac{1}{\sqrt{2}} \frac{d\sigma(\mathcal{M}'_d = 1) + d\sigma(\mathcal{M}'_d = -1) - 2d\sigma(\mathcal{M}'_d = 0)}{d\sigma(\mathcal{M}'_d = 1) + d\sigma(\mathcal{M}'_d = -1) + d\sigma(\mathcal{M}'_d = 0)}, \quad (10)$$

is predicted to be $1/\sqrt{2}$ from the selection rules. Indeed, the full evaluation of Eqs. (6-8) yields a value close to that (see Fig. 4).

Apart from the polarization observables which near the threshold are determined directly by the corresponding quantities for the subprocess $pn \rightarrow d\phi$, one can define another set of observables with completely different behavior in the two reactions, i.e., in the subprocess $pn \rightarrow d\phi$ and in the reaction (1). For this purpose let us define observables at fixed initial deuteron and proton spin projections, e.g., at $\mathcal{M}_d = 0$ and $s_p = \frac{1}{2}$, respectively. Near the threshold this means that the spin projection of the neutron will be $s_n = -s'_p$ and by measuring the spectator polarization one can form different combinations which determine (allowed or forbidden) transitions for the subprocess $pn \rightarrow d\phi$. Together with conservation rules for the complete reaction this admits a study of different observables which vanish in the sub-reaction $pn \rightarrow d\phi$. Correspondingly, let us define the quantity

$$W_{zz} = \left. \frac{d\sigma(\mathcal{M}_\phi = 1) - d\sigma(\mathcal{M}_\phi = -1)}{d\sigma(\mathcal{M}_\phi = 1) + d\sigma(\mathcal{M}_\phi = -1)} \right|_{s_p=\frac{1}{2}; \mathcal{M}_d=0}. \quad (11)$$

If one measures this quantity for the positive spin projection of the spectator, then in the subprocess $pn \rightarrow d\phi$ one has super allowed transitions ($s_p + s_n = 0$, $\mathcal{M}_\phi = \pm 1$, $\mathcal{M}'_d = \mp 1$) and, since in (11) a summation over the final deuteron projections is assumed, W_{zz} is predicted to approach zero at the threshold (see [16]).

A different situation occurs if one considers W_{zz} at $s'_p = -1/2$. In this case, at least for $t = t_{min(max)}$ (i.e., the forward/backward direction for the subprocess $pn \rightarrow d\phi$), the conservation of the spin projection in the complete reaction (1) strongly suppresses the final

states with $\mathcal{M}_\phi = \mathcal{M}'_d = \pm 1$ and $\mathcal{M}_\phi = -1, \mathcal{M}'_d = 0$, so that the only contributing final states remain these with $\mathcal{M}_\phi = 1, \mathcal{M}'_d = 0, -1$ and $\mathcal{M}_\phi = -1, \mathcal{M}'_d = 1$, which imply $W_{zz} \sim 1/3$, in contrast to the vanishing W_{zz} in $pn \rightarrow d\phi$ [16]. Figure 5 illustrates our full numerical evaluation for W_{zz} at $s'_p = -1/2$.

IV. FACTORIZATION

Let us discuss in more detail the behavior of the cross section Eq. (2) with the matrix element (8) near the threshold. As mentioned above, the summation over the spin projection of the intrinsic neutron s_n in Eq. (8) prevents an exact factorization of the cross section into the cross section of the subprocess $pn \rightarrow d\phi$ and a “target factor”. However, from Figs. 3 - 5 we conclude that, since the deviations of the exactly calculated observables from those determined by the selection rules of the subprocess $pn \rightarrow d\phi$ are small, a factorization should hold indeed with a good accuracy. For instance, the asymmetry in Fig. 3 demonstrates the validity of the selection rule $s_p = -s_n$, i.e., the amplitude $T_{s_p, s_n}^{\mathcal{M}_\phi, \mathcal{M}'_d}(\nu_{pn}, t)$ is proportional to $\delta_{s_p, -s_n}$. Similarly, Fig. 5 shows that the main contribution comes from transitions with $|\mathcal{M}'_d| = 1$, and from the above selection rules the amplitude $T_{s_p, s_n}^{\mathcal{M}_\phi, \mathcal{M}'_d}(\nu_{pn}, t)$ is also proportional to $\delta_{\mathcal{M}'_d, -\mathcal{M}_\phi}$. Hence, near the threshold one may represent the amplitude as $T_{s_p, s_n}^{\mathcal{M}_\phi, \mathcal{M}'_d}(\nu_{pn}, t) = \delta_{\mathcal{M}'_d, -\mathcal{M}_\phi} \delta_{s_p, -s_n} \tilde{T}_{s_p, \mathcal{M}'_d}(\nu_{pn}, t)$, where $\tilde{T}_{s_p, \mathcal{M}'_d}(\nu_{pn}, t)$ is the threshold-near amplitude of the process $pn \rightarrow d\phi$. Then in Eq. (8) the summation over s_n disappears and, as conjectured above, the unpolarized cross section Eq. (2) factorizes. Substituting this amplitude into Eq. (2), introducing the corresponding flux factors for the sub-reaction $pn \rightarrow d\phi$ and performing the summation over all spin projections, the unpolarized cross section can be cast in the form

$$2E'_d \frac{d^5\sigma}{d^3p'_d d\Omega'_p} = \frac{2}{(2\pi)^3} \frac{\nu_{pn}(\nu_{pn} - 4m^2) |\mathbf{p}'_p|^2}{|\mathbf{p}| |B|\mathbf{p}'_p| - CE'_p} \left(u_S^2(|\mathbf{p}'_p|) + u_D^2(|\mathbf{p}'_p|) \right) \frac{d\sigma(pn \rightarrow d\phi)}{dt}, \quad (12)$$

where the differential cross section $d\sigma(pn \rightarrow d\phi)/dt$ depends only upon the transferred momentum t (determined by p_p and p'_d) and the invariant mass ν_{pn} (determined by p_p and p'_p due to the balance $E_n = M_d - E'_p$, $\mathbf{p}_n = -\mathbf{p}'_p$). From Eq. (12) it is seen that by studying experimentally the amplitude (8), near the threshold, one can obtain directly information about the subprocess $pn \rightarrow d\phi$. Here it is worth reminding that the factorization of the cross section holds only if the contribution of the D wave is negligible, i.e., if in Eq. (8) one can put $s_n = -s_p$ and arrives at

$$\mathcal{F} \approx \sqrt{8\pi M_d} T_{s_p, -s_p}^{\mathcal{M}_\phi, \mathcal{M}'_d}(\nu_{pn}, t) \left[u_S(|\mathbf{p}'_p|) C_{\frac{1}{2}s'_p \frac{1}{2}-s_p}^{1\mathcal{M}'_d} - u_D(|\mathbf{p}'_p|) \sqrt{4\pi} \sum_{m\mu} C_{\frac{1}{2}s'_p \frac{1}{2}-s_p}^{1\mu} C_{2m1\mu}^{1\mathcal{M}'_d} Y_{2m}(\mathbf{n}) \right]. \quad (13)$$

Eqs. (8, 12) and (13) show that the smaller the values of the spectator momentum are the higher the accuracy of the factorization is. This has been confirmed numerically by comparing exact results based on Eqs. (2, 8) and approximate results based on Eqs. (12, 13). In the whole kinematical region of interests the difference is less than 0.5%.

One may generally conjecture that a structure analog to Eq. (8) gives rise to factorization under suitable kinematical conditions, hence providing a theoretical foundation of quasi-free reactions at the neutron in the deuteron.

V. SUMMARY

In summary, within a covariant Bethe-Salpeter approach we study the relation of the tagged quasi-free reaction $pn \rightarrow d\phi$ and the complete process $pd \rightarrow d\phi p_{sp}$ near the threshold. Guided by the very proximity of the numerical results for cross section and polarization observables based on the exact formulae and on the factorized ones, we find that, for special kinematical conditions, selection rules are operative immediately causing a factorization $\sigma(pd \rightarrow d\phi p_{sp}) = F \sigma(pn \rightarrow d\phi)$ where F depends on deuteron and kinematical quantities. This is equivalent to an omission of the contribution of the deuteron D wave at small values of the spectator momentum.

As an extension of the present work, an investigation of the twice tagged reaction $dd \rightarrow nnMp_{sp1}p_{sp2}$ can be envisaged to study the quasi-free meson production by the nn reaction proposed in [19].

ACKNOWLEDGMENTS

We are grateful to A.I. Titov for many valuable discussions. L.P.K. and S.S.S. acknowledge the warm hospitality of the nuclear theory group in the Research Center Rossendorf. The work has been supported by Heisenberg - Landau program, BMBF grant 06DR921, and RFBR 00-15-96737.

REFERENCES

- [1] T. Peterson et al., Nucl. Phys. A **663**, 1057c (2000).
- [2] P. Moskal, M. Wolke, A. Khoukaz, W. Oelert, Prog. Part. Nucl. Phys. **49**, 1 (2002).
- [3] A.I. Titov, B. Kämpfer, V.V. Shklyar, Phys. Rev. C **59**, 999 (1999);
A.I. Titov, B. Kämpfer, B.L. Reznik, Phys. Rev. C **65**, 065202 (2002).
- [4] T. Chiavassa et al., Phys. Lett. B **337**, 192 (1994).
- [5] H. Calén et al., Phys. Rev. Lett. **79**, 2642 (1997), **80** (1998) 2069, Phys. Rev. C **58**, 2667 (1998);
R. Bilger et al., Nucl. Phys. A **663**, 1053c (2000); Nucl. Instr. Meth. A **457**, 64 (2001).
- [6] M. Büscher et al., "Study of ω and ϕ meson production in the reaction $pd \rightarrow dVp_{sp}$ at ANKE", experiment proposal #75/ANKE, <http://ikpd15.ikp.kfa-juelich.de:8085/doc/Proposals.html>;
www.fz-juelich.de/ikp/anke/doc/Proposals.shtml.
- [7] L.A. Kondratyuk, Ye.S. Golubeva, M. Büscher, nucl-th/9808050;
V.Yu. Grishina, L.A. Kondratyuk, M. Büscher, nucl-th/9906064;
V.Yu. Grishina, L.A. Kondratyuk, M. Büscher, J. Haidenbauer, C. Hanhart, J. Speth, Phys. Lett. B **475**, 9 (2000).
- [8] K. Nakayama, J. Haidenbauer, J. Speth, Nucl. Phys. A **689**, 402c (2001); Phys. Rev. C **63**, 015201 (2000).
- [9] A. Sibirtsev, W. Cassing, Eur. Phys. J. A **7**, 407 (2000).
- [10] A.I. Titov, M. Fujiwara, T.S.H. Lee, Phys. Rev. C **66**, 022202 (2002);
A.I. Titov, T.S.H. Lee, Phys. Rev. C **66**, 015204 (2002);
Y. Oh, A.I. Titov, T.S.H. Lee, Phys. Rev. C **63**, 025201 (2001).
- [11] A.I. Titov, B. Kämpfer, B.L. Reznik, Eur. Phys. J. A **7**, 543 (2000).
- [12] F. Balestra (DISTO Collaboration), Phys. Rev. C **63**, 024004 (2001);
B. Kämpfer, R. Kotte, C. Hartnack, J. Aichelin, J. Phys. G **28**, 2035 (2002).
- [13] J. Friese (for the HADES Collaboration), Prog. Part. Nucl. Phys. **42**, 235 (1999).
- [14] A.I. Titov, B. Kämpfer, Eur. Phys. J. A **12**, 217 (2001).
- [15] G. Penner, U. Mosel, Phys. Rev. C **65**, 055202 (2002);
K. Tsushiba, K. Nakayama, nucl-th/0211065.
- [16] L.P. Kaptari, B. Kämpfer, Eur. Phys. J. A **14**, 211 (2002).
- [17] L.P. Kaptari, A.Yu. Umnikov, S.G. Bondarenko, K.Yu. Kazakov, F.C. Khanna, B. Kämpfer, Phys. Rev. C **54**, 986 (1996).
- [18] A.Yu. Umnikov, L.P. Kaptari, F.C. Khanna, Phys. Rev. C **56**, 1700 (1997);
A.Yu. Umnikov, L.P. Kaptari, K.Yu. Kazakov, F.C. Khanna, Phys. Lett. B **334**, 163 (1994);
A.Yu. Umnikov, Z. Phys. A **357**, 333 (1997).
- [19] P. Moskal et al., Phys. Lett. B **517**, 295 (2001).

FIGURES




FIG. 1. Diagram of the process (1).




FIG. 2. Differential cross section of the reaction $pd \rightarrow d\phi p_{sp}$ as a function of the deuteron momentum p'_d at $p_p = 3.505 \text{ GeV}/c$, $\theta_{d'} = 0^\circ$, $\theta_{p'_{sp}} = 130^\circ$.




FIG. 3. As Fig. 2 but for the asymmetry A .




FIG. 4. As Fig. 2 but for T_{20} .




FIG. 5. As Fig. 2 but for W_{zz} .